

UR-1529
ER/40685/917

ALTERNATIVES TO AXION PRODUCTION AND DETECTION*

by

A.C. Melissinos
Department of Physics and Astronomy
University of Rochester
Rochester, NY 14627, USA

*To appear in Proceedings of the 5th IFT Workshop on Axions, University of Florida, Gainesville, FL, March 13-15, 1998.

1. Heavy Ion Collisions

Axions are associated with the Peccei-Quinn symmetry which is broken at very high energies ($f \gtrsim 10^{10}$ GeV). However axions acquire mass near the QCD phase transition ($E \sim 1$ GeV). Thus it should be possible to produce axions in heavy ion collisions (at RHIC and the LHC) where the equivalent “temperature” of the quark-gluon plasma will exceed the 1 GeV mark. We estimate the expected rate of axion production and detection as follows [1].

The number density of axions at time t_1 , where $T(t_1) = T_1 \sim 1$ GeV is given by

$$\begin{aligned} n_a(t_1) &\simeq \rho_a(t_1) \frac{1}{m_a(t_1)} \\ &\simeq \frac{1}{2} v^2 m_a(t_1) < \alpha^2(t_1) > \\ &\simeq \pi f_a^2 \frac{1}{t_1} = \frac{1}{2} f_a^2 m_a \end{aligned} \tag{1}$$

For the variables used in Eq. (1) we note that v and α are related to the vacuum expectation value of the scalar field $\phi(x)$

$$< \phi(x) > = v e^{i\alpha(x)}$$

and thus the axion field is given by

$$a(x) = v\alpha(x)$$

We have also introduced the axion coupling $f_a = v/N$ where N is the “number of minima” in the axion effective potential, and $< \alpha(x) > \sim 1/N$. Furthermore

$$f_a m_a \simeq f_\pi m_\pi$$

where f_π , m_π are the pion decay constant and mass, for ~ 0.1 GeV, $m_\pi \sim 0.14$ GeV. Finally we can take

$$t_1 m_a(t_1) \simeq 2\pi$$

Thus we expect the axion density in a quark-gluon plasma at $T \sim 1$ GeV to be ($1 \text{ eV} = 5 \times 10^4 \text{ cm}^{-3}$)

$$\begin{aligned} n_a &= \frac{1}{2}(f_\pi m_\pi)^2 \frac{1}{m_a} \simeq 10^{37} \left(\frac{10^{-5} \text{ eV}}{m_a} \right) \text{ eV}^3 \\ &\simeq 10^{12} \left(\frac{10^{-5} \text{ eV}}{m_a} \right) \text{ F}^{-3} \end{aligned} \quad (2)$$

To detect the axions produced we will use their conversion to photons in a magnetic field [2]. The conversion probability is given by

$$P_{a \rightarrow \gamma} = \frac{1}{4} g_{a\gamma\gamma}^2 (B\ell)^2 \quad (3)$$

with B , ℓ the magnetic field strength and length and $g_{a\gamma\gamma}$ the axion to 2-photon coupling

$$g_{a\gamma\gamma} = \frac{\alpha}{2\pi} \frac{N}{f_a} \quad (4)$$

Combining Eqs. (1,3 and 4) we find the probability for detection of a photon from axion conversion

$$P_\gamma = \left(\frac{\alpha}{2\pi} \right)^2 \frac{N^2}{8} (B\ell)^2 m_a V \quad (5)$$

where V is the volume of the quark gluon plasma region. It is interesting that Eq. (5) depends on the axion field parameters linearly and only through the axion mass m_a .

We evaluate Eq. (5) for typical parameters $B = 5 \text{ T} = 10^3 \text{ eV}^2$,

$$\ell = 10 \text{ m} = 5 \times 10^7 \text{ eV}^{-1},$$

$$V \simeq (10 \text{ F})^3 = 10^{-22} \text{ eV}^{-3}$$

and set

$$(\alpha/2\pi)^2 (N^2/8) \simeq 10^{-6}$$

to find

$$P_\gamma = 2.5 \times 10^{-7} \left(\frac{m_a}{1\text{eV}} \right) / \text{collision} \quad (6)$$

Assuming single bunches in RHIC and one central collision per beam crossing (5×10^4 collisions/second) the expected photon rate is

$$P_\gamma \sim 0.7 \left(\frac{m_a}{1\text{eV}} \right) / \text{minute} \quad (7)$$

In writing Eq. (5) we assumed that all axions produced during the quark-gluon formation period cross the magnet and reach the detector. Such a condition can be approached by exploiting the Lorentz contraction for asymmetric collisions. If A is the area and L the distance of the detector from the collision point, we must have

$$1/\gamma_{\text{cm}}^2 \geq A/L^2 \quad (8)$$

For typical values $A \sim 100 \text{ cm}^2$, $L = 10 \text{ m}$ one finds $\gamma > 100$. This is easy to achieve in a fixed target experiment but in a collider it will be necessary to optimize the trade-off between cm energy and detector acceptance.

Finally we estimate the energy of the detected photons. Presumably in the cm system the axion energy will be representative of the plasma temperature, $E_a \sim 1 \text{ GeV}$. If the collision is asymmetric the additional boost results in higher energies at the detector. Thus one is searching for high energy photons (multi-GeV) which appear to have traversed through the shielding surrounding the collision point. Such a signal is unambiguous, especially since it depends quadratically on the value of the converting magnetic field.

2. Critical Field

It has been recently shown that in a critical electromagnetic field, e^+e^- pairs are spontaneously produced [3, 4]. The critical field is defined [5] by

$$E_c = \frac{m^2 c^3}{e \hbar} = 1.3 \times 10^{16} \text{ V/cm} \quad (9)$$

and diagrammatically, the pair production can be viewed as the multiphoton process shown in Fig. 1a. The pair can

couple to produce an axion as shown in Fig. 1b, if the e - e - a coupling is different from zero.

The probability per unit time and unit volume for graph (a) and for strong fields is given by [6]

$$w_{e^+e^-} = \frac{\alpha E^2}{\pi \hbar^2} \exp\left(-\frac{\pi m^2 c^3}{e \hbar E}\right) \quad (10)$$

For graph (b) we multiply Eq. (10) by the e - e - a coupling squared and by an inverse area typical of the interaction region. Since the e^+e^- pair is produced at one space-time point within a volume of typical dimensions of the electron's Compton wavelength, we write

$$w_a = \frac{\alpha E^2}{\pi \hbar^2} \frac{m_e^2 c^2}{\hbar^2} \left(\frac{\hbar c}{f_a}\right)^2 \exp\left(-\frac{\pi m^2 c^3}{e \hbar E}\right) \quad (11)$$

For $E \simeq \pi E_c$ the number of axions produced in volume-time VT is given by

$$N_a \simeq \left(\frac{m_e}{f_a}\right)^2 \frac{V c T}{\lambda_c^4} \quad (12)$$

Near-critical fields in the laboratory frame should be reached by focusing to the diffraction limit the beam from the next generation of 1Å X-ray Free Electron Lasers [7]. We then take $V = (10\text{\AA})^3$ and $T = 0.2$ ps with a repetition frequency of 7×10^4 Hz (for the DESY FEL), to obtain an axion

rate

$$R_a = 5 \times 10^{-9} \left(\frac{m_a}{\text{eV}} \right)^2 \text{ s}^{-1} \quad (13)$$

This is too low a rate for considering laboratory experiments since the axion detection efficiency is equally small.

Axion production in a critical field could, however, be important in astrophysical settings where the magnetic field of neutron stars can reach critical value. If this is the case, axions with an energy $E_a \sim 1$ MeV should be emitted. If we assume that V in Eq. (12) is given by the entire volume of the neutron star ($R = 10$ km) located at a distance of 0.3 kpc, then we obtain for the flux on earth

$$F_a \simeq 5 \left(\frac{m_a}{\text{eV}} \right)^2 \text{ m}^{-2} \text{ s}^{-1} \quad (14)$$

Still an excessively low rate, but the process could be of interest if a proper calculation reveals the possibility of higher fluxes.

References

- [1] P. Sikivie in “Proceedings of the Workshop on Cosmic Axions”, p233 World Scientific 1990, C. Jones and A. Melissinos eds.
- [2] G. Ruoso *et al*, Z. für Physik C **56**, 505 (1992).
- [3] D.L. Burke *et al*, Phys. Rev. Lett. **79**, 1626 (1997).
- [4] A.C. Melissinos, “The spontaneous breakdown of the vacuum”, to appear in proceedings of the Conference on “Quantum Aspects of Particle Beams”, World Scientific 1998. Also University of Rochester report UR-1520.
- [5] J. Schwinger, Phys. Rev. **82**, 664 (1951); **93**, 615 (1954).
- [6] E. Brezin and C. Itzykson, Phys. Rev. **D2**, 1191 (1970).
- [7] See for instance “A VUV Free Electron Laser at the TESLA Facility at DESY”, DESY print June 1995 and “LCLS Design Study Report”, SLAC-R-1521, 1998. Available from NTIS, Springfield VA, 22161.

This figure "ur1529fig.GIF" is available in "GIF" format from:

<http://arXiv.org/ps/hep-ph/9805506v1>